



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

inspected at the International foot and carriage bridge, 3,225; persons inspected at the International foot and carriage bridge refused entry, 9.

Statistical reports of States and cities of the United States—Yearly and monthly.

CALIFORNIA—*Los Angeles*.—Month of September, 1904. Estimated population, 160,000. Total number of deaths, 225, including diphtheria 3, enteric fever 5, and 45 from tuberculosis.

Stockton.—Month of September, 1904. Estimated population, 20,000. Total number of deaths, 22, including 3 from tuberculosis.

COLORADO—*Colorado Springs*.—Month of September, 1904. Estimated population, 30,000. Total number of deaths, 34, including enteric fever 3, and 12 from tuberculosis.

CONNECTICUT—*Bridgeport*.—Month of September, 1904. Estimated population, 82,128. Total number of deaths, 104, including enteric fever 3, and 13 from tuberculosis.

LOUISIANA—*New Orleans*.—Month of September, 1904. Estimated population, 317,000; white, 233,000; colored, 84,000. Total number of deaths, 471; white, 295; colored, 176, including diphtheria 1, enteric fever 12, whooping cough 2, and 96 from tuberculosis.

MASSACHUSETTS—*Brockton*.—Month of September, 1904. Estimated population, 46,601. Total number of deaths, 40, including 5 from phthisis pulmonalis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended October 8, 1904, from 87 observers, indicate that scarlet fever, smallpox, inflammation of brain, and puerperal fever were more than usually prevalent, and phthisis pulmonalis, dysentery, cholera morbus, intermittent fever, cholera infantum, pleuritis, remittent fever, erysipelas, pneumonia, whooping cough, measles, and meningitis were less than usually prevalent.

Whooping cough was reported present at 5, measles at 11, pneumonia at 14, diphtheria at 30, smallpox at 42, scarlet fever at 48, enteric fever at 117, and phthisis pulmonalis at 295 places.

MINNESOTA—*Winona*.—Month of September, 1904. Estimated population, 23,000. Total number of deaths, 23, including diphtheria 1 and 2 from phthisis pulmonalis.

NEW YORK—*Auburn*.—Month of September, 1904. Estimated population, 38,500. Total number of deaths, 24, including diphtheria 2, enteric fever 1, and 5 from tuberculosis.

NORTH CAROLINA.—Reports to the State board of health for the month of August, 1904, from 22 towns, having an aggregate population of 142,950 (white, 83,350; colored, 59,600), show a total of 272 deaths—white, 116; colored, 156 (including enteric fever 27, whooping cough 1, and 34 from phthisis pulmonalis).

OHIO.—Reports to the State board of health for the nine weeks ended September 3, 1904, from 64 localities, having an aggregate population of 1,284,134, show 23 deaths from diphtheria, 66 from enteric fever, 14 from measles, 6 from scarlet fever, and 5 from whooping cough.

Cleveland.—Month of September, 1904. Estimated population, 420,000. Total number of deaths, 478, including diphtheria 6, enteric fever 11, and 45 from tuberculosis.

Columbus.—Month of September, 1904. Census population, 125,560. Total number of deaths, 161, including enteric fever 6, scarlet fever 1, and 21 from tuberculosis.

East Liverpool.—Month of August, 1904. Estimated population, 20,000. Total number of deaths, 25, including diphtheria 1, and 3 from tuberculosis.

Month of September, 1904. Total number of deaths, 15, including whooping cough 1 and 2 from phthisis pulmonalis.

PENNSYLVANIA—*Altoona*.—Month of September, 1904. Census population, 38,973. Total number of deaths, 53, including diphtheria 2, enteric fever 3, whooping cough 1, and 5 from tuberculosis.

TENNESSEE—*Nashville*.—Month of September, 1904. Estimated population, 82,947—white, 52,061; colored, 30,886. Total number of deaths, 146—white, 69; colored, 77, including enteric fever 9, scarlet fever 1, whooping cough 2, and 22 from tuberculosis.

UTAH—*Ogden*.—Month of September, 1904. Estimated population, 20,000. Total number of deaths, 24, including diphtheria 1, enteric fever 1, and 1 from scarlet fever.

Salt Lake City.—Month of September, 1904. Census population, 53,531. Total number of deaths, 65, including diphtheria 1, enteric fever 7, and 5 from tuberculosis.

VIRGINIA—*Richmond*.—Month of September, 1904. Estimated population, 100,000; white, 62,250; colored, 37,750. Total number of deaths, 144; white 68, colored 76, including enteric fever 6, and 17 from tuberculosis.